



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

except that the captives seemed to be the better favored. If one were inclined to conscientious scruples against keeping wild birds in confinement the sight of this happy and contented family would certainly sweep the scruples away. We feel a lasting gratitude to Mr. Worthington for his interest and liberality in making this venture possible, and we heartily congratulate Dr. Scott upon the beginning of a realization of the fruition of a life given to a greatly needed work such as he has undertaken. May his work prosper and his days be multiplied to carry forward this work.

Carefully prepared lists of summer birds from regions not already fully covered by published lists, are solicited. Faunal studies of this sort will be worth publishing until the country is fully covered, or until we have learned everything there is to learn about distribution. Along with any list there should go such notes relating to life histories as can be learned, particularly as to the close of the season of song, and exact dates of nesting. Copious notes made while one is working in the field with the birds are valuable assets. Often what seem to be purposeless notes at the time they are made later prove to be the key to the solution of some vexed question. The best memory may prove fickle.

Before the next number of the Bulletin goes to press the southward migrations will have begun. How much do we really know about when the southward movement begins in our own locality, or where we spend the summer? Isn't it worth while to begin to learn about it? Here on Lake Erie some of the sandpipers which nest about Hudson's Bay, supposably, are to be found on the sandy shores and in the mud flats early in July, and some of the warblers are evidently moving southward early in August. On the other hand, some birds seem to leave for the south before they actually do.

SOME NOTEWORTHY LORAIN COUNTY RECORDS FOR 1906.

A solitary White-throated Sparrow was found wintering in the gorge of Vermillion river near Brownhelm Mills, on January 1. The bird was clearly in good physical condition. This is the first recorded instance of the wintering of this species.

Hoyt's Horned Lark was again found, in company with Prairie Horned Larks, four individuals on February 22. The difference of coloration and size was marked. These four inclined to separate themselves from the others and grouped together.

A company of six Bronzed Grackles spent the winter in Oberlin. Two or more Robins were fed all winter in Oberlin.

Only two Lapland Longspurs have been recorded for the county—March 5. Many more are usually found during the spring weeks.

A solitary Snowflake was found feeding with a company of Prairie Horned Larks, on February 16. No others were seen during the winter.

A Kirtland's and a fine male Prairie Warbler were recorded on May 2. The birds were feeding close together and permitted an approach within twenty feet, singing all the time. The Kirtland's was without markings on the center of the breast. This is the third record for Kirtland's Warbler.

A solitary Hooded and a solitary Kentucky Warbler were recorded on May 15, both singing lustily. These birds have recently made their appearance in the county. Neither have yet been found except during the height of the spring migrations.

There was a notable scarcity of Goldfinches during the last week in March and the first three weeks in April. Only a few were recorded during the winter. The birds became common again by the 30th of April, and before any had completed the spring moult. A similar scarcity is reported to me from south-eastern Michigan. Were they common anywhere during this period?

SPECIAL PAPERS RECEIVED.

Cassinia, A Bird Annual. Proceedings of the Delaware Valley Ornithological Club of Philadelphia. 1905. Issued February, 1906.

On a Collection of Birds and Mammals from the Colorado Delta, Lower California. By Witmer Stone, with Field Notes by Samuel N. Rhoads. From the Proceedings of The Academy of Natural Sciences of Philadelphia, September, 1905. Issued December 6, 1905.

On a Collection of Birds from British East Africa obtained by Mr. George L. Harrison, Jr. By Witmer Stone. From the Proceedings of The Academy of Natural Sciences of Philadelphia, November, 1905. Issued January 24, 1906.

The Mammals of Colorado. By Edward R. Warren. Colorado College Publications, General Series No. 19.

CURRENT PUBLICATIONS RECEIVED.

American Ornithology, Vol. V, No. 12; Vol. VI, Nos. 1 to 5.

Bird-Lore, Vol. VII, No. 6; Vol. VIII, Nos. 1, 2.

Boys and Girls, Vol. V, Nos. 3 to 6; Vol. VI, Nos. 1 to 5.